

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/99234/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

El-Hiti, Gamal A., Smith, Keith, Alamri, Mesfer, Morris, Ceri A., Kariuki, Benson ORCID: <https://orcid.org/0000-0002-8658-3897> and Kille, Peter ORCID: <https://orcid.org/0000-0001-6023-5221> 2017. Crystal structure of 2-(bis(4-methoxyphenyl)amino)-2-oxoacetic acid, C₁₆H₁₅NO₅. Zeitschrift für Kristallographie - New Crystal Structures 232 (2) 10.1515/ncrs-2016-0325 file

Publishers page: <http://dx.doi.org/10.1515/ncrs-2016-0325>
<<http://dx.doi.org/10.1515/ncrs-2016-0325>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See


<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Open Access

Gamal A. El-Hiti*, Keith Smith, Mesfer Alamri, Ceri A. Morris, Benson M. Kariuki and Peter Kille

Crystal structure of 2-(bis(4-methoxyphenyl)amino)-2-oxoacetic acid, $C_{16}H_{15}NO_5$



DOI 10.1515/ncrs-2016-0325

Received October 17, 2016; accepted January 3, 2017; available online January 20, 2017

Abstract

$C_{16}H_{15}NO_5$, monoclinic, $P2_1/n$ (no. 14), $a = 6.7689(5)$ Å, $b = 45.219(3)$ Å, $c = 10.1102(6)$ Å, $\beta = 101.360(7)^\circ$, $V = 3033.9(4)$ Å³, $T = 298(2)$ K.

CCDC no.: 1525292

Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

A solution of oxalyl chloride (1 mole equivalent) in dichloromethane (DCM) was added dropwise to a solution of bis(4-methoxyphenyl)amine (1 mole equivalent) in DCM in the presence of triethylamine at room temperature. The mixture was stirred for 1.5 h and water was added. The organic layer was separated, dried over anhydrous

Table 1: Data collection and handling.

Crystal:	Colorless block
Size:	$0.30 \times 0.22 \times 0.17$ mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	0.10 mm^{-1}
Diffractometer, scan mode:	Atlas, ω
$2\theta_{\text{max}}$, completeness:	58° , 97.3%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	15585, 6846, 0.353
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3929
$N(\text{param})_{\text{refined}}$:	403
Programs:	CrysAlis ^{PRO} [18], SHELX [19], WinGX [20], CHEMDRAW Ultra [21]

magnesium sulfate and evaporated under reduced pressure to give the title compound in 46% yield. The low yield could be a result of half of the amine acting as a base to abstract hydrogen chloride evolved from the reaction. To investigate this issue the reaction was repeated with two equivalents of bis(4-methoxyphenyl)amine and no triethylamine. Following aqueous work-up, the crude product was obtained in 75% yield based on oxalyl chloride. Crystallization using acetonitrile gave the title compound as colorless crystals, Mp. 122–123 °C. The NMR spectra recorded at room temperature showed two sets of signals for the two aryl rings, confirming restricted rotation about the C–N bond. The barriers to free rotation in such compounds are already known to be substantial [1–3]. ¹H NMR (400 MHz, DMSO- d_6): δ 7.04, 6.96 (2 d, $J = 8.5$ Hz, 4 H, H-2/H-6), 6.86, 6.69 (2 d, $J = 8.5$ Hz, 4 H, H-3/H-5), 3.82, 3.71 (2 s, 6 H, OMe); ¹³C NMR (100 MHz, DMSO- d_6): δ 164.4 (s, CO₂H), 159.2 (s, C=O), 158.3 (s, C-4), 134.0, 132.8 (2 s, C-1), 129.8, 127.7 (2 d, C-2/C-6), 114.8 (2 d, C-3/C-5), 56.0, 55.8 (2 q, OMe); ES⁺ – MS: m/z (%) 302 (MH⁺, 100), 288 (12), 256 (21), 228 (12); HRMS (ES⁺): calculated for $C_{16}H_{16}NO_5$ (MH⁺): 302.1028; found: 302.1028. IR (FT): ν_{max} 3300, 1740, 1713, 1665, 1500, 1366, 1167 cm^{−1}.

Experimental details

Non-hydrogen atoms were refined with anisotropic displacement parameters. All hydrogen atoms were placed in calculated positions and refined using a ring model. Methyl

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Mesfer Alamri: School of Biosciences, Cardiff University, Cardiff CF10 3AT, UK; and School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Ceri A. Morris: School of Biosciences, Cardiff University, Cardiff CF10 3AT, UK; and School of Medicine, Cardiff University, Tenovus Building, Heath Park, Cardiff CF14 4XN, UK

Peter Kille: School of Biosciences, Cardiff University, Cardiff CF10 3AT, UK

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	U_{iso}^*/U_{eq}
C1	−0.3411(7)	0.26406(10)	−0.1012(4)	0.1293(17)
H1A	−0.4716	0.2549	−0.1178	0.194*
H1B	−0.3171	0.2728	−0.1831	0.194*
H1C	−0.3359	0.2791	−0.0336	0.194*
C2	−0.2196(4)	0.22459(6)	0.0489(3)	0.0588(7)
C3	−0.0928(4)	0.20057(6)	0.0757(3)	0.0633(7)
H3	0.0066	0.1975	0.0254	0.076*
C4	−0.1137(4)	0.18125(6)	0.1767(3)	0.0610(7)
H4	−0.0282	0.1650	0.1943	0.073*
C5	−0.2587(4)	0.18557(5)	0.2517(2)	0.0477(6)
C6	−0.3827(5)	0.20947(7)	0.2264(3)	0.0724(9)
H6	−0.4807	0.2126	0.2777	0.087*
C7	−0.3637(5)	0.22914(7)	0.1245(3)	0.0789(9)
H7	−0.4489	0.2454	0.1076	0.095*
C8	−0.3962(4)	0.14275(6)	0.3527(2)	0.0507(6)
C9	−0.5151(4)	0.13514(6)	0.2117(3)	0.0504(6)
C10	−0.1313(4)	0.17106(5)	0.4890(2)	0.0496(6)
C11	0.0482(4)	0.15607(6)	0.5151(2)	0.0576(7)
H11	0.0779	0.1424	0.4531	0.069*
C12	0.1853(4)	0.16125(6)	0.6335(3)	0.0629(7)
H12	0.3077	0.1512	0.6510	0.075*
C13	0.1395(4)	0.18131(6)	0.7255(3)	0.0596(7)
C14	−0.0412(5)	0.19644(6)	0.6983(3)	0.0633(7)
H14	−0.0713	0.2101	0.7601	0.076*
C15	−0.1772(4)	0.19128(6)	0.5798(3)	0.0595(7)
H15	−0.2992	0.2014	0.5616	0.071*
C16	0.4445(5)	0.17124(8)	0.8805(3)	0.0927(11)
H16A	0.4118	0.1506	0.8745	0.139*
H16B	0.5101	0.1759	0.9711	0.139*
H16C	0.5328	0.1758	0.8198	0.139*
C17	1.5746(4)	0.04762(7)	1.5043(3)	0.0717(8)
H17A	1.5792	0.0303	1.4501	0.108*
H17B	1.6307	0.0431	1.5969	0.108*
H17C	1.6512	0.0632	1.4738	0.108*
C18	1.2664(4)	0.06381(5)	1.3672(2)	0.0468(6)

C—H bonds were fixed at 0.96 Å and displacement parameters were 1.5 times $U_{eq}(C)$. The methyl groups were allowed to spin about the C—C bond. Aromatic C—H distances were set to 0.93 Å and their $U(iso)$ parameters were set to 1.2 times $U_{eq}(C)$. Hydroxyl O—H distances were set to 0.82 Å and their $U(iso)$ set to 1.5 times $U_{eq}(O)$. Crystal data, data collection and structure refinement details are summarized in Table 1.

Comment

Aryl oxamic acid derivatives have various interesting applications [4–8]. In addition, aryl oxamic acids can be used as intermediates for the synthesis of various classes of compounds including heterocycles [9–12]. Oxamates can be synthesized by the use of various synthetic procedures [13–17].

In the title crystal structure, the asymmetric unit consists of two independent molecules of C₁₆H₁₅NO₅. The oxoacetic acid fragments of the molecule are involved in intermolecular hydrogen bonding, of the type O—H...O, with the following geometric parameters: O4...O7 = 2.673(2) Å, O—H...O = 171.0°; and O9...O2 = 2.734(2) Å, O—H...O = 169.9° forming chains along [101].

These hydrogen bonds can be classified as medium strong. Bond lengths and angles in both crystallographically independent molecules are in the expected ranges.

Acknowledgements: We thank the EPSRC for the grant which supplied the MS instrumentation used in this study. M. Alamri thanks the Saudi Cultural Bureau, London for a scholarship and G. A. El-Hiti extends his appreciation to the Deanship of Scientific Research at King Saud University for its funding for this research through the research group project RGP-239.

References

- Hobson, R. F.; Reeves, L. W.: Hindered rotation about the N-C bond in some vinylogous amides. *J. Magn. Reson.* **10** (1973) 243–252.
- Smith, B. D.; Goodenough-Lashua, D. M.; D'Souza, C. J. E.; Norton, K. J.; Schmidt, L. M.; Tung, J. C.: Substituent effects on the barrier to carbamate C–N rotation. *Tetrahedron Lett.* **45** (2004) 2747–2749.
- Krishnan, V. V.; Thompson, W. B.; Goto, J. J.; Maitra, K.; Maitra, S.: Modulations in restricted amide rotation by steric induced conformational trapping. *Chem. Phys. Lett.* **523** (2012) 124–127.
- Maiore, L.; Aragoni, M. C.; Carcangiu, G.; Cocco, O.; Isaia, F.; Lippolis, V.; Meloni, P.; Murru, A.; Slawin, A. M. Z.; Tuveri, E.; Woollins, J. D.; Arca, M.: Oxamate salts as novel agents for the restoration of marble and limestone substrates: case study of ammonium *N*-phenyloxamate. *New J. Chem.* **40** (2016) 2768–2774.
- Miskimins, W. K.; Ahn, H. J.; Kim, J. Y.; Ryu, S.; Jung, Y.-S.; Choi, J. Y.: Synergistic anti-cancer effect of phenformin and oxamate. *PLOS One* **9** (2014) e85576, doi: 10.1371/journal.pone.0085576.
- Choi, S.-R.; Beeler, A. B.; Pradhan, A.; Watkins, E. B.; Rimoldi, J. M.; Tekwani, B.; Avery, M. A.: Generation of oxamic acid libraries: antimalarials and inhibitors of *plasmodium falciparum* lactate dehydrogenase. *J. Comb. Chem.* **9** (2007) 292–300.
- Hargrave, K. D.; Hess, F. K.; Oliver, J. T.: *N*-(4-Substituted-thiazolyl)oxamic acid derivatives, new series of potent, orally active antiallergy agents. *J. Med. Chem.* **26** (1983) 1158–1163.
- Klaubert, D. H.; Sellstedt, J. H.; Guinasso, C. J.; Capetola, R. J.; Bell, S. C.: *N*-(Aminophenyl)oxamic acids and esters as potent, orally active antiallergy agents. *J. Med. Chem.* **24** (1981) 742–748.

9. Wang, H.; Guo, L.-N.; Wang, S.; Duan, X.-H.: Decarboxylative alkynylation of α -keto acids and oxamic acids in aqueous media. *Org. Lett.* **17** (2015) 3054–3057.
10. Loloiu, G.; Maior, O.: Isatin chemistry. Synthesis of *N*-methyl-2,3-dioxo-2,3-dihydropyrrolo(2,3-*b*) phenoxathiin. *Rev. Roum. Chim.* **42** (1997) 67–69.
11. Molina, P.; Vilaplana, M. J.; Andreu, P. L.; Moller, J.: Oxamic acid derivatives in heterocyclic synthesis: preparation of 1,2,4-triazolo[1,5-*a*]pyrazine derivatives. *J. Heterocycl. Chem.* **24** (1984) 1281–1284.
12. Downs, J. R.; Pastine, S. J.; Schady, D. A.; Greer, H. A.; Kelley, W.; Embree, M. C.; Townsend, J. D.; Beam, C. F.: Preparation of 1*H*-pyrazole-5-carboxamides from dilithiated C(α),*N*-phenylhydrazones and lithiated ethyl oxanilates or lithiated ethyl oxamate. *J. Heterocycl. Chem.* **38** (2001) 691–694.
13. Gadge, S. T.; Kusumawati, E. N.; Harada, K.; Sasaki, T.; Nishio-Hamane, D.; Bhanage, B. M.: Synthesis of oxamate and urea by oxidative single and double carbonylation of amines using immobilized palladium metal-containing ionic liquid@SBA-15. *J. Mol. Catal. A: Chem.* **400** (2015) 170–178.
14. Gadge, S. T.; Bhanage, B. M.: Pd/C-Catalyzed synthesis of oxamates by oxidative cross double carbonylation of amines and alcohols under Co-catalyst, base, dehydrating agent, and ligand-free conditions. *J. Org. Chem.* **78** (2013) 6793–6797.
15. Lisnard, L.; Chamoreau, L.-M.; Li, Y.; Journaux, Y.: Solvothermal synthesis of oxamate-based helicate: temperature dependence of the hydrogen bond structuring in the solid. *Cryst. Growth Des.* **12** (2012) 4955–4962.
16. Yang, G.; Zhang, H.; Huang, Y.; Chen, Z.: Synthesis of methyl *N*-aryl oxamate using soluble polymer support. *Synth. Commun.* **36** (2006) 611–619.
17. Lesimple, P.; Bigg, D. C. H.: An improved procedure for the preparation of alkyl *N*-(4-aryl-2-thiazolyl)oxamates. *Synthesis* **1991**(9) (1991) 763–764.
18. Agilent. CrysAlis^{PRO}. Agilent Technologies, Yarnton, England, 2014.
19. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
20. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
21. Cambridge Soft. CHEMDRAW Ultra. Cambridge Soft Corporation, Cambridge, MA, USA, 2001.